
Name of the Student:_____
Max. Marks : 26 Marks **Time : 26 Minutes**


Q1.

Figure 1 shows apparatus used to investigate the inverse-square law for gamma radiation.

Figure 1

Downloaded from www.merit-minds.com





A sealed source that emits gamma radiation is held in a socket attached to clamp **B**. The vertical distance between the open end of the source and the bench is 138 mm. A radiation detector, positioned vertically above the source, is attached to clamp **T**.

A student is told **not** to move the stands closer together.

(a) Describe a procedure for the student to find the value of d, the vertical distance between the open end of the source and the radiation detector.

In your answer, annotate above the figure to show how a set-square can be used in this procedure.

Downloaded from www.merit-minds.com

_____

_____

_____

_____

_____

(2)

(b) Before the source was brought into the room, a background count C_b was recorded.

C_b = 630 counts in 15 minutes

With the source and detector in the positions shown in the figure above, d = 530 mm. Separate counts C_1, C_2 and C_3 are recorded.

C_1 = 90 counts in 100 s
C_2 = 117 counts in 100 s
C_3 = 102 counts in 100 s

R_C is the mean count rate corrected for background radiation.

Show that when d = 530 mm, R_C is about 0.3 s^{-1}.

(2)

(c) The apparatus is adjusted so that d = 380 mm.
Counts are made that show R_C = 0.76 s^{-1}.

The student predicts that:

$$R_C = \frac{k}{d^2}$$

where k is a constant.

Explain whether the values of R_C in parts (b) and (c) support the student's prediction.

Downloaded from www.merit-minds.com


(2)

(d) Describe a safe procedure to reduce d. Give a reason for your procedure.

(2)

The student determines R_C for further values of d.
The values of d change by the same amount Δd between each measurement.
Figure 2 shows these data.

Figure 2



(e) Determine Δd.

Downloaded from www.merit-minds.com

Δd = _____ mm

(2)

(f) Explain how the student could confirm whether the graph above supports the prediction:

$$R_C = \frac{k}{d^2}$$

No calculation is required.

_____

_____

_____

_____

_____

_____

_____

_____

_____

(3)

When a gamma photon is detected by the detector, another photon cannot be detected for a time t_d called the dead time.

It can be shown that:

$$t_d = \frac{R_2 - R_1}{R_1 \times R_2}$$

where R_1 is the measured count rate

R_2 is the count rate when R_1 is corrected for dead time error.

(g) The distance between the source and the detector is adjusted so that d is very small and R_1 is 100 s^{-1}.
On average, two of the gamma photons that enter the detector every second are not detected.

Calculate t_d for this detector.

Downloaded from www.merit-minds.com

t_d = _____ s

(1)

(h) A student says that if 100 gamma photons enter a detector in one second and t_d is 0.01 s, all the photons should be detected.

Explain, with reference to the nature of radioactive decay, why this idea is **not** correct.

_____

_____

_____

_____

_____


_____

(2)

(Total 16 marks)

Q2.

The diagram shows alpha particles all travelling in the same direction at the same speed. The alpha particles are scattered by a gold $(^{197}_{79}\text{Au})$ nucleus.
The path of alpha particle **1** is shown.



(a) State the fundamental force involved when alpha particle **1** is scattered by the nucleus in the diagram.

_____

(1)

(b) Draw an arrow at position **X** on the diagram above to show the direction of the rate of change in

Downloaded from www.merit-minds.com

momentum of alpha particle **1**

(1)

(c) Suggest **one** of the alpha particles in the diagram above which may be deflected downwards with a scattering angle of 90°

Justify your answer.

alpha particle number = _____

_____

_____

_____

_____

(2)

(d) Alpha particle **4** comes to rest at a distance of 5.5×10^{-14} m from the centre of the $^{197}_{79}Au$ nucleus.

Calculate the speed of alpha particle **4** when it is at a large distance from the nucleus. Ignore relativistic effects.

mass of alpha particle = 6.8×10^{-27} kg

speed = _____ m s^{-1}

(3)

(e) The nuclear radius of $^{197}_{79}Au$ is 6.98×10^{-15} m.

Calculate the nuclear radius of $^{107}_{47}Ag$.

Downloaded from www.merit-minds.com

radius = _____ m

(2)

(f) All nuclei have approximately the same density.

State **one** conclusion about the nucleons in a nucleus that can be deduced from this fact.

_____

_____

_____

_____

(1)

(Total 10 marks)

Downloaded from www.merit-minds.com