
Name of the Student:_____
Max. Marks : 24 Marks **Time : 24 Minutes**

Q1.

Titration curves, labelled **E**, **F**, **G** and **H**, for combinations of different aqueous solutions of acids and bases are shown below.

All solutions have concentrations of 0.1 mol dm^{-3}.



(a) In this part of the question, write the appropriate letter in each box.

From the curves **E**, **F**, **G** and **H**, choose the curve produced by the addition of

(i) sodium hydroxide to 25 cm^3 of ethanoic acid ☐

(1)

(ii) ammonia to 25 cm^3 hydrobromic acid ☐

Downloaded from www.merit-minds.com

(1)

(iii) hydrochloric acid to 25 cm^3 of potassium hydroxide ☐

(1)

(b) The table shows information about some acid-base indicators.

Indicator	pH range	Lower pH colour	Higher pH colour
pentamethoxy red	1.2−3.2	violet	colourless
naphthyl red	3.7−5.0	red	yellow
4−nitrophenol	5.6−7.0	colourless	yellow
cresol purple	7.6−9.2	yellow	purple

(i) Which indicator in the table could be used for the titration that produces curve **E** but **not** for the titration that produces curve **F**?

Tick (✓) **one** box.

pentamethoxy red ☐

naphthyl red ☐

4−nitrophenol ☐

cresol purple ☐

(1)

(ii) Give the colour change at the end point of the titration that produces curve **H** when naphthyl red is used as the indicator.

_____

(1)

(iii) A beaker contains 25 cm^3 of a buffer solution at pH = 6.0

Two drops of each of the four indicators in the table are added to this solution.

State the colour of the mixture of indicators in this buffer solution.

You should assume that the indicators do **not** react with each other.

_____

(1)

(Total 6 marks)

Downloaded from www.merit-minds.com

Q2.

Water dissociates slightly according to the equation:

$$H_2O(l) \rightleftharpoons H^+(aq) + OH^-(aq)$$

The ionic product of water, K_w, is given by the expression

$$K_w = [H^+][OH^-]$$

K_w varies with temperature as shown in the table.

Temperature / °C	**K_w / mol^2 dm^{-6}**
25	1.00×10^{-14}
50	5.48×10^{-14}

(a) Explain why the expression for K_w does **not** include the concentration of water.

(2)

(b) Explain why the value of K_w increases as the temperature increases.

(2)

(c) Calculate the pH of pure water at 50 °C.
Give your answer to 2 decimal places.

Downloaded from www.merit-minds.com

(3)

(d) Calculate the pH of 0.12 mol dm^{-3} aqueous NaOH at 50 °C.
Give your answer to 2 decimal places.

(3)
(Total 10 marks)

Downloaded from www.merit-minds.com

Q3.

The acid dissociation constant, K_a, for ethanoic acid is given by the expression

$$K_a = \frac{[H^+][CH_3COO^-]}{[CH_3COOH]}$$

The value of K_a for ethanoic acid is 1.74×10^{-5} mol dm^{-3} at 25 °C.

(a) A buffer solution is prepared using ethanoic acid and sodium ethanoate. In the buffer solution, the concentration of ethanoic acid is 0.186 mol dm^{-3} and the concentration of sodium ethanoate is 0.105 mol dm^{-3}.

Calculate the pH of this buffer solution.
Give your answer to 2 decimal places.

(3)

(b) In a different buffer solution, the concentration of ethanoic acid is 0.251 mol dm^{-3} and the concentration of sodium ethanoate is 0.140 mol dm^{-3}.

A sample of hydrochloric acid containing 0.015 mol of HCl is added to 1000 cm^3 of this buffer solution.

Calculate the pH of the buffer solution after the hydrochloric acid has been added.
You should ignore any change in total volume.
Give your answer to 2 decimal places.

Downloaded from www.merit-minds.com

_____

_____

_____

(5)

(Total 8 marks)

Downloaded from www.merit-minds.com