Name of the Student:_____
Max. Marks : 16 Marks **Time : 16 Minutes**

Q1.

The value of the acid dissociation constant, K_a, for ethanoic acid is 1.74×10^{-5} mol dm^{-3} at 298 K.

(a) (i) Write an expression for K_a for ethanoic acid.

_____

(ii) Calculate the pH at 298 K of a 0.220 mol dm^{-3} solution of ethanoic acid.

_____

_____

_____

_____

_____

_____

(5)

(b) A sample of the 0.220 mol dm^{-3} solution of ethanoic acid was titrated against sodium hydroxide solution.

(i) Calculate the volume of a 0.150 mol dm^{-3} solution of sodium hydroxide required to neutralise 25.0 cm^3 of the ethanoic acid solution.

_____

_____

_____

_____

(ii) From the list below, select the best indicator for this titration and explain your choice.

Name of indicator	**pH range**
bromophenol blue	3.0 – 4.6
methyl red	4.2 – 6.3
bromothymol blue	6.0 – 7.6
thymol blue	8.0 – 9.6

Indicator _____

Explanation _____

Downloaded from www.merit-minds.com

_____

_____

(5)

(c) A buffer solution is formed when 2.00 g of sodium hydroxide are added to 1.00 dm^3 of a 0.220 mol dm^{-3} solution of ethanoic acid.

Calculate the pH at 298 K of this buffer solution.

_____

_____

_____

_____

_____

_____

_____

(6)

(Total 16 marks)

Downloaded from www.merit-minds.com