

Name of the Student: \_\_\_\_\_

Max. Marks : 25 Marks

Time : 25 Minutes

Q1.

Aspirin can be produced by reacting salicylic acid with ethanoic anhydride.  
An incomplete method to determine the yield of aspirin is shown.

1. Add about 6 g of salicylic acid to a weighing boat.
2. Place the weighing boat on a 2 decimal place balance and record the mass.
3. Tip the salicylic acid into a 100 cm<sup>3</sup> conical flask.
4. \_\_\_\_\_
5. Add 10 cm<sup>3</sup> of ethanoic anhydride to the conical flask and swirl.
6. Add 5 drops of concentrated phosphoric acid.
7. Warm the flask for 20 minutes.
8. Add ice-cold water to the reaction mixture and place the flask in an ice bath.
9. Filter off the crude aspirin from the mixture and leave it to dry.
10. Weigh the crude aspirin and calculate the yield.

- (a) Describe the instruction that is missing from step 4 of the method.

Justify why this step is necessary.

Instruction \_\_\_\_\_

Justification \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_

(2)

- (b) Suggest a suitable piece of apparatus to measure out the ethanoic anhydride in step 5.

\_\_\_\_\_


(1)

- (c) Identify a hazard of using concentrated phosphoric acid in step 6.

\_\_\_\_\_

(1)

- (d) Complete the equation for the reaction of salicylic acid with ethanoic anhydride to produce aspirin.



(1)

- (e) A 6.01 g sample of salicylic acid ( $M_r = 138.0$ ) is reacted with  $10.5 \text{ cm}^3$  of ethanoic anhydride ( $M_r = 102.0$ ). In the reaction the yield of aspirin is 84.1%

The density of ethanoic anhydride is  $1.08 \text{ g cm}^{-3}$

Show by calculation which reagent is in excess.

Calculate the mass, in g, of aspirin ( $M_r = 180.0$ ) produced.

Reagent in excess \_\_\_\_\_

Mass of aspirin \_\_\_\_\_ g

(5)

- (f) Suggest **two** ways in which the melting point of the crude aspirin collected in step 9 would differ from the melting point of pure aspirin.

Difference 1 \_\_\_\_\_

\_\_\_\_\_

Difference 2 \_\_\_\_\_

\_\_\_\_\_

(2)

- (g) The crude aspirin can be purified by recrystallisation using hot ethanol (boiling point =  $78^\circ\text{C}$ ) as the solvent.

Describe **two** important precautions when heating the mixture of ethanol and crude aspirin.

Precaution 1 \_\_\_\_\_

\_\_\_\_\_

Precaution 2 \_\_\_\_\_

\_\_\_\_\_

(2)

- (h) The pure aspirin is filtered under reduced pressure.  
A small amount of cold ethanol is then poured through the Buchner funnel.

Explain the purpose of adding a small amount of cold ethanol.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

(1)

- (i) A sample of the crude aspirin is kept to compare with the purified aspirin.

Describe **one** difference in appearance you would expect to see between these two solid samples.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

(1)

(Total 16 marks)

Q2.

This question is about citric acid, a hydrated tricarboxylic acid. Its formula can be represented as  $\text{H}_3\text{Y} \cdot x\text{H}_2\text{O}$

- (a) A 1.50 g sample of  $\text{H}_3\text{Y} \cdot x\text{H}_2\text{O}$  contains 0.913 g of oxygen by mass.  
The sample burns completely in air to form 1.89 g of  $\text{CO}_2$  and 0.643 g of  $\text{H}_2\text{O}$

Show that the empirical formula of citric acid is  $\text{C}_3\text{H}_5\text{O}_4$


(5)

- (b) A 3.00 g sample of  $\text{H}_3\text{Y} \cdot x\text{H}_2\text{O}$  ( $M_r = 210.0$ ) is heated to constant mass. The anhydrous  $\text{H}_3\text{Y}$  that remains has a mass of 2.74 g

Show, using these data, that the value of  $x = 1$

(2)

The figure shows the structure of  $\text{H}_3\text{Y}$



- (c) Complete this IUPAC name for  $\text{H}_3\text{Y}$

\_\_\_\_\_ propane-1, 2, 3-tricarboxylic acid

(1)

- (d) State the number of peaks you would expect in the  $^{13}\text{C}$  NMR spectrum for  $\text{H}_3\text{Y}$

