

Name of the Student: _____

Max. Marks : 17 Marks

Time : 17 Minutes

Q1.

- (a) Use data from the table below to explain why dilute hydrochloric acid cannot be used to acidify potassium manganate(VII) in a titration.

	$E^\ominus / V$
$MnO_4^-(aq) + 8H^+(aq) + 5e^- \rightarrow Mn^{2+}(aq) + 4H_2O(l)$	+1.51
$Cl_2(aq) + 2e^- \rightarrow 2Cl^-(aq)$	+1.36
$2H^+(aq) + 2e^- \rightarrow H_2(aq)$	0.00

(2)

- (b) Use information from the table in part (a) to determine the minimum volume, in cm^3 , of $0.500 \text{ mol dm}^{-3}$ sulfuric acid that is required for a titre of 25.0 cm^3 of $0.0200 \text{ mol dm}^{-3}$ potassium manganate(VII) solution.
Show your working.

(3)

- (c) In each titration using potassium manganate(VII), a large excess of dilute sulfuric acid is used to avoid any possibility of the brown solid MnO_2 forming.

- (i) Deduce a half-equation for the reduction of MnO_4^- ions in acidic solution to form MnO_2 .

(1)

- (ii) Give **two** reasons why it is essential to avoid this reaction in a titration between potassium manganate(VII) and iron(II) ions.

(2)

- (d) Potassium manganate(VII) is an oxidising agent.
Suggest **one** reason why a $0.0200 \text{ mol dm}^{-3}$ solution of potassium manganate(VII) does **not** need to be kept away from flammable material.

(1)

(Total 9 marks)

Q2.

Some electrode potentials are shown in the table below. These values are **not** listed in numerical order.

Electrode half-equation	$E^\ominus / \text{V}$
$\text{Cl}_2(\text{aq}) + 2\text{e}^- \longrightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$2\text{HOCl}(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \longrightarrow \text{Cl}_2(\text{aq}) + 2\text{H}_2\text{O}(\text{l})$	+1.64
$\text{H}_2\text{O}_2(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \longrightarrow 2\text{H}_2\text{O}(\text{l})$	+1.77
$\text{O}_2(\text{g}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \longrightarrow \text{H}_2\text{O}_2(\text{aq})$	+0.68
$\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \longrightarrow 2\text{H}_2\text{O}(\text{l})$	+1.23

- (a) Identify the most powerful reducing agent from all the species in the table.

(1)

- (b) Use data from the table to explain why chlorine should undergo a redox reaction with water. Write an equation for this reaction.

Explanation _____

Equation _____

(2)

- (c) Suggest **one** reason why the redox reaction between chlorine and water does not normally occur in the absence of light.

(1)

- (d) Use the appropriate half-equation from the table to explain in terms of oxidation states what happens to hydrogen peroxide when it is reduced.

(2)

- (e) Use data from the table to explain why one molecule of hydrogen peroxide can oxidise another molecule of hydrogen peroxide. Write an equation for the redox reaction that occurs.

Explanation _____

Equation _____

(2)

(Total 8 marks)