Name of the Student:_____
Max. Marks : 21 Marks **Time : 21 Minutes**

Mark Schemes

Q1.

(a) moles of $Cr_2O_7^{2-}$ per titration = 21.3 × 0.0150 / 1000 = $\underline{3.195 \times 10^{-4}}$

1

($Cr_2O_7^{2-}$ + 14H^+ + 6Fe^{2+} → 2Cr^{3+} + 7H_2O + 6Fe^{3+}) $Cr_2O_7^{2-}$:Fe^{2+} = 1:6

If 1:6 ratio incorrect cannot score M2 or M3

1

moles of Fe^{2+} = 6 × 3.195 × 10^{-4} = 1.917 × 10^{-3}

Process mark for M1 × 6 (also score M2)

1

original moles in 250 cm^3 = 1.917 × 10^{-3} × 10 = 1.917 × 10^{-2}

Process mark for M3 × 10

1

mass of $FeSO_4.7H_2O$ = 1.917 × 10^{-2} × 277.9 = 5.33 (g)

Mark for answer to M4 × 277.9

(allow 5.30 to 5.40)

*Answer **must** be to at least 3 sig figs*

Note that an answer of 0.888 scores M1, M4 and M5 (ratio 1:1 used)

1

(b) (Impurity is a) reducing agent / reacts with dichromate / impurity is a version of $FeSO_4$ with fewer than 7 waters (not fully hydrated)

Allow a reducing agent or compound that that converts Fe^{3+} into Fe^{2+}

1

Such that for a given mass, the impurity would react with more dichromate than a similar mass of $FeSO_4.7H_2O$

OR for equal masses of the impurity and $FeSO_4.7H_2O$, the impurity would react with more dichromate.

Must compare mass of impurity with mass of $FeSO_4.7H_2O$

1

[7]

Downloaded from www.merit-minds.com

Q2.

(a) Hydrogen/H_2 gas/bubbles

1

1.0 mol dm^{-3} HCl/H^+

1

At 298K and 100kPa

Allow 1 bar instead of 100 kPa
Do not allow 1 atm

1

Pt (electrode)

1

(b) $Li^+ + MnO_2 + e^- \rightarrow LiMnO_2$

Ignore state symbols

1

–0.13(V)

1

(c) Fe^{3+} ions reduced to Fe^{2+}

Can score from equation/scheme

1

Because $E(Fe^{3+}/Fe^{2+}) > E(H^+/H_2)$/$E$(hydrogen)

Allow emf/E_{cell} +ve/0.77V
Allow Fe^{3+} better oxidising agent than H^+
Allow H_2 better reducing agent than Fe^{2+}
Only award this explanation mark if previous mark given

1

(d) Moles $Cr_2O_7^{2-}$ = $\underline{23.7 \times 0.01/1000}$ = 2.37×10^{-4}

1

1 mol $Cr_2O_7^{2-}$ reacts with 6 mol Fe^{2+} so moles Fe^{2+} in 25 cm^3 = $6 \times 2.37 \times 10^{-4} = 1.422 \times 10^{-3}$

1

M1 × 6

Moles Fe^{2+} in 250 cm^3 = 1.422×10^{-2}

M2 × 10 or M4/10

1

Original moles Fe^{2+} = $\underline{10.00/277.9}$ = 0.0360

Independent mark

1

Moles Fe^{2+} oxidised = 0.0360 – 0.0142 = 0.0218

M4 – M3

1

% oxidised = (0.0218 × 100)/0.0360 = 60.5%

(M5 × 100)/M4
Allow 60 to 61
Note Max 3 if mol ratio for M2 wrong
eg 1:5 gives 67.1%
1:1 gives 93.4%
Note also, 39.5% (39-40) scores M1, M2, M3 and M4 (4 marks)

Downloaded from www.merit-minds.com

[14]

Downloaded from www.merit-minds.com